

## MATH 1420: WORKSHEET FOR SECTION 3.2

There are two types of rules for derivatives: rules that tell you how to differentiate a basic function, and rules that tell you what to do when a function is built up from simpler functions. Here's some rules to get us started.

### RULES FOR BASIC FUNCTIONS

**Constant Rule.** Whenever  $c$  is a constant,

$$\frac{d}{dx}c = 0.$$

**Power Rule.** When  $n$  is a positive integer

$$\frac{d}{dx}x^n = nx^{n-1}.$$

**Extended Power Rule.** More generally this works for any (nonzero) power:

$$\frac{d}{dx}x^\alpha = \alpha x^{\alpha-1}.$$

### RULES FOR BUILDING UP FROM SIMPLER FUNCTIONS

**Multiplication by a constant.**

$$\frac{d}{dx}cf(x) = c\frac{d}{dx}f(x) = cf'(x).$$

**Adding functions.**

$$\frac{d}{dx}(f(x) + g(x)) = \frac{d}{dx}f(x) + \frac{d}{dx}g(x) = f'(x) + g'(x).$$

**Product rule.**

$$\frac{d}{dx}f(x)g(x) = f'(x)g(x) + f(x)g'(x).$$

**Quotient rule.**

$$\frac{d}{dx}\frac{f(x)}{g(x)} = \frac{f'(x)g(x) - f(x)g'(x)}{g(x)^2}$$

Using these rules, you can compute the derivative of any polynomial or any rational function. Here's some functions to differentiate.

(1)  $a(x) = x^2 = x \cdot x$ . Try this with both the power rule and the product rule.

(2)  $b(x) = 4x^5 - 3x^2 + 2x + 3$ .

(3)  $c(x) = \frac{1}{x}$ .

(4)  $d(x) = \frac{1}{x^3}$ . Try this with both the extended power rule and the quotient rule.

(5)  $e(x) = \frac{(x-1)(x+1)}{x^2}$ .

(6)  $f(x) = 1 - \frac{1}{x^2}$ . (You should get a formula for  $f'(x)$  equivalent to what you got for  $e'(x)$ . Why?)

(7)  $g(x) = 3x^2 + \frac{2x-1}{x+2}$ .