

Math 195: Logarithms

Julia Williams

Cape Cod Community College

Spring 2026

Last time

- Last time we saw **exponential functions**, those of the form

$$f(x) = a^x, \quad \text{where } a \neq 1 \text{ is positive.}$$

- Today we'll look at a new kind of function, **logarithms**, which are closely related to exponential functions.

What is a logarithm?

- A **logarithm** is an inverse of an exponential function.
- Write $\log_a(x)$ for the inverse of a^x . We call a the **base** of the logarithm.
- Like with exponential functions, a must be positive and cannot be 1. Usually we only look at $a > 1$.

Special notation

- $\ln x = \log_e x$ is the natural logarithm.

Special notation

- $\ln x = \log_e x$ is the **natural logarithm**.
- The textbook writes $\log x$ for $\log_{10} x$, sometimes called the **common logarithm**.
- **Caution!** Different sources are inconsistent about what $\log x$ means.
 - Mathematicians and most programming languages mean \ln .
 - Computer scientists sometimes mean \log_2 .
 - Other sources use it to mean \log_{10} .

My advice: never write $\log x$, always explicitly write the base.

What are logarithms saying?

$$\log_a x = y \iff a^y = x$$

This means that computing $\log_a x$ is asking, how many copies of a do you multiply together to get x ?

What are logarithms saying?

$$\log_a x = y \iff a^y = x$$

This means that computing $\log_a x$ is asking, how many copies of a do you multiply together to get x ?

- $\log_{10} 1000 = 3$ because $10^3 = 1000$.
- $\log_2 128 = 7$ because $2^7 = 128$.
- $\log_4 2 = \frac{1}{2}$ because $\sqrt{4} = 2$.
- $\log_3(\frac{1}{9}) = -2$ because $3^{-2} = \frac{1}{9}$.

What are logarithms saying?

$$\log_a x = y \iff a^y = x$$

This means that computing $\log_a x$ is asking, how many copies of a do you multiply together to get x ?

- $\log_{10} 1000 = 3$ because $10^3 = 1000$.
- $\log_2 128 = 7$ because $2^7 = 128$.
- $\log_4 2 = \frac{1}{2}$ because $\sqrt{4} = 2$.
- $\log_3(\frac{1}{9}) = -2$ because $3^{-2} = \frac{1}{9}$.

This is not very useful when x is not a neat power of a !

- $\ln 5 = \text{????}$

Almost always, $\log_a x$ is an **irrational number**.

Graphs of logarithms, and global behavior

Knowing that logarithms are inverses of exponential functions tells us how their graphs look.

Graphs of logarithms, and global behavior

Knowing that logarithms are inverses of exponential functions tells us how their graphs look.

$\log_a x$, where $a > 1$ has the following global behavior:

- $\text{dom}(\log_a) = (0, \infty)$
- $\text{ran}(\log_a) = \mathbb{R}$
- \log_a is increasing on its entire domain.
- \log_a is concave down on its entire domain.

Geometric transformations of logarithms

Like with other functions, knowing the graph of $\log_a x$ lets us understand graphs of its geometric transformations.

- For logarithms, the most important feature of the graph is the vertical asymptote. Its location is determined by the horizontal shift, if any.
- The other important feature is its orientation. Which reflections were done to the mother function?

A little bit of history

- The invention of the logarithm was one of the most important mathematical discoveries in history.

A little bit of history

- The invention of the logarithm was one of the most important mathematical discoveries in history.
- First published by John Napier in 1614.
- *Mirifici Logarithmorum Canonis Descriptio* contained tables of values of logarithms of trigonometric functions of different inputs.
- Logarithms were a tool to make calculations easier.

$$\log_a(xy) = \log_a x + \log_a y \quad \log_a(x^n) = n \log_a x$$

- This use only became obsolete with the rise of the electronic digital computer.
- While they are no longer an essential tool for arithmetic, logarithms continue to have many important uses.



John Napier, 1616